



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

where the following abbreviations are used:

$$\begin{aligned} a &= a_1a_2, \quad b = b_1b_2, \quad A = a_2 - a_1, \quad B = b_2 - b_1, \\ C &= a_1b_2 - a_2b_1, \quad D = a_1b_2 + a_2b_1, \quad L = C[ADm + (m+2)aB], \\ M &= C[BDm + (m+2)bA], \quad N = a(C^2m + aB^2), \quad P = b(C^2m + bA^2), \\ Q &= maAC, \quad R = mbBC, \quad S = (mC^2D + 2abAB). \end{aligned}$$

If we look upon (8) as an algebraical equation in  $p = dy/dx$  which has 4 roots  $p_1, p_2, p_3, p_4$ , these being functions of  $x$  and  $y$ , then we are led to the conclusion that through any point in the plane there are *four* directions satisfying the condition proposed. We have, therefore, *four* curves in question.

For the present purpose the equation (8) in  $p$  may be considered irreducible. Then, if a singular solution of this equation exists, it must simultaneously satisfy the equations

$$\varphi = 0, \quad \frac{\partial \varphi}{\partial p} = 0, \quad \frac{\partial \varphi}{\partial x} + p \frac{\partial \varphi}{\partial y} = 0.$$

The last equation is *identically zero* in the present case, and we have

$$\begin{aligned} \frac{d\varphi}{dp} &= 4p^3Qx - 3p^2(C^2x^2 - Lx + Py + N) - 2p(2C^2xy \\ &\quad + Mx - Ly - S) - (C^2y^2 + My - Rx + P) = 0. \end{aligned} \quad (9)$$

Equations (8) and (9), therefore, represent, in parametric form, the singular solution of the differential equation (8), which is, then, the algebraic curve in question.

#### 411. Proposed by C. N. SCHMALL, New York City.

$ABCD$  is a rectangle of known sides.  $BC$  being produced indefinitely, it is required to draw a straight line from  $A$  cutting  $CD$  and  $BC$  in  $X$  and  $Y$ , respectively, so that the intercept  $XY$  may be equal to a given straight line. (Unsolved in *Educational Times*.)

### II. SOLUTION BY THE PROPOSER.

We shall assume that the given rectangle is a square, the problem thus being a special case of problem 382, proposed by R. C. Archibald in the May (1911) number of the MONTHLY.

CONSTRUCTION: Along  $AB$  produced lay off  $AE$  equal to the *given length*. Draw  $ED$ . Prolong  $AD$  to  $K$  so that  $DK = DE$ . On  $AK$  as a diameter (centre  $O$ ) describe a semi-circle cutting  $BC$  produced in  $Y$ . Draw  $AY$  cutting  $DC$  in  $X$ . Then  $AXY$  is the line required; *i. e.*, the intercept  $XY$  is of the given length.

PROOF: Draw  $YH$  perpendicular to  $AK$ . Draw  $XK$  and  $YK$ . Then the right triangles  $ADX$  and  $YHK$  are equal in all respects (congruent). For the angles  $DAX$  and  $HYK$  are equal, and  $DA = DC = HY$ .

